

# NIR Excitation and Tunneling of Dihydroxycarbene



**Project Manager**  
Prof. Dr. Peter R. Schreiner

**Researchers**  
Bastian Bernhardt, Henrik Quanz,  
Frederik R. Erb, Marcus A. Bartlett,  
Wesley D. Allen and Prof. Dr. Peter R.  
Schreiner

**Principal Investigator**  
Bastian Bernhardt

**Project Term**  
2017 - 2019

**Project Areas**  
Molecular Chemistry

**Clusters**  
Goethe-HLR Frankfurt, FUCHS-CSC  
Frankfurt

**Software**  
GAUSSIAN

**Additional Software**  
CFOUR

**Institute**  
Institut für Organische Chemie

**University**  
Justus Liebig University Giessen

**Partners**  
Center for Computational Quantum  
Chemistry, University of Georgia

## Introduction

While quantum-mechanical tunneling (QMT) was recognized early in the emerging field of quantum theory as an accelerating effect in chemical reactions,[1] its fundamental importance beyond kinetic and thermodynamic control for chemical reactions was demonstrated only recently.[2,3] As a means to an end, matrix isolation was utilized to generate, trap, and identify reactive intermediates that undergo spontaneous QMT.[4] In case of higher-energy conformers, selective excitation with near-infrared light (NIR) is suitable to prepare one desired conformer.[5–8] Such experiments provide insights into medium effects[9] and conformation dependent QMT.[10,11] In the project presented herein, we investigated dihydroxycarbene by experimental and computational means. Dihydroxycarbene is an activated isomer of formic acid,[12] that displays an intricate QMT process.

## Methods

We use the matrix isolation technique combined with IR spectroscopy to characterize the three conformers of dihydroxycarbene (**tt**, **ct**, and **cc**; **c** = *cis*, **t** = *trans*), which are generated through pyrolysis of oxalic acid. The trapped conformers can be interconverted into each other by irradiating the matrix with an optical parametric oscillator at a given wavelength. Simultaneously, we compute the corresponding potential energy surface (PES) of this system at the coupled cluster level of theory. The computations are also crucial to compare experimental with predicted IR spectra and tunneling half-lives.

## Results

*Cis-cis*-dihydroxycarbene (**cc**) was prepared for the first time in a nitrogen matrix at 3 K; this is the first *cis-cis*-hydroxycarbene observed to date. NIR excitation of **tt** induces the reaction **tt** → **ct** + **cc**, while excitation of **ct** yields **tt** and **cc**. The latter conformer then tunnels with a half-life of ca. 15 min back to **ct**. This value is in excellent agreement with our computations, that predicted a half-life of 18 min. Experiments in argon did not result in the formation of **cc** during NIR irradiation but only the conversion **tt** → **ct** and its backreaction were observed. Furthermore, we were able to correct earlier reported PES for this system significantly by applying a focal point scheme. At this level, the classical activation barrier for the interconversion **cc** → **ct** is 9.3 kcal mol<sup>-1</sup> which is about 1.5 kcal mol<sup>-1</sup> lower than predicted by computations at a lower level of theory. As QMT is very sensitive towards the barrier width and height, these computations increase our understanding about the reactivity of dihydroxycarbene enormously.

## Discussion

Our experiments suggest that the half-life of **cc** in argon is very small. It is somewhat surprising that our gas-phase computations seem to be more comparable to the results in N<sub>2</sub> than in Ar. The NIR induced process for the dideuterated species has not yet been successful in either matrix material as the activation barrier lies above the excitation energy in this case. However, irradiation of a combination band of **d<sub>2</sub>-tt** at 5257 cm<sup>-1</sup> (15.03 kcal mol<sup>-1</sup>) over 3 h resulted in a very small increase of **d<sub>2</sub>-ct** bands in argon. If this observation is due to vibrationally activated tunneling,[13] the background radiation of the OPO (355 nm), or just an artifact of the difference spectrum, will be investigated in further experiments.

## Outlook

In order to obtain experimental isotope shifts for **cc** and compare them to computed values, experiments on the monodeuterated isotopologues will be performed. We have already been able to compute the PES and tunneling half-lives for this system. In future studies the *cis-trans*-interconversions of other hydroxycarbenes will be investigated in order to get a deeper insight into tunneling processes of such reactive intermediates. We are continuing to exploit both experimental and computational methods for this purpose.

## Publications

## Reference

- [1] Bell, R. P. (1936): The theory of reactions involving proton transfers. Proceedings of the Royal Society of London. Series A-Mathematical and Physical Sciences, 154(882), 414-429. <https://doi.org/10.1098/rspa.1936.0060>
- [2] Schreiner, P. R., Reisenauer, H. P., Ley, D., Gerbig, D., Wu, C. H., & Allen, W. D. (2011): Methylhydroxycarbene: Tunneling control of a chemical reaction. Science, 332(6035), 1300-1303. <https://doi.org/10.1126/science.1203761>
- [3] Schreiner, P. R. (2017): Tunneling control of chemical reactions: the third reactivity paradigm. Journal of the American Chemical Society, 139(43), 15276-15283. <https://doi.org/10.1021/jacs.7b06035>
- [4] Pettersson, M., Lundell, J., Khriachtchev, L., & Räsänen, M. (1997): IR spectrum of the other rotamer of formic acid, cis-HCOOH. Journal of the American Chemical Society, 119(48), 11715-11716. <https://doi.org/10.1021/ja972362i>
- [5] Maçôas, E. M., Khriachtchev, L., Pettersson, M., Fausto, R., & Räsänen, M. (2003): Rotational isomerism in acetic acid: the first experimental observation of the high-energy conformer. Journal of the American Chemical Society, 125(52), 16188-16189. <https://doi.org/10.1021/ja038341a>
- [6] Lapinski, L., Reva, I., Rostkowska, H., Halasa, A., Fausto, R., & Nowak, M. J. (2013). Conformational transformation in squaric acid induced by near-IR laser light. The Journal of Physical Chemistry A, 117(25), 5251-5259. <https://doi.org/10.1021/jp402128g>
- [7] Ryazantsev, S. V., Feldman, V. I., & Khriachtchev, L. (2017): Conformational Switching of HOCO Radical: Selective Vibrational Excitation and Hydrogen-Atom Tunneling. Journal of the American Chemical Society, 139(28), 9551-9557. <https://doi.org/10.1021/jacs.7b02605>
- [8] Jesus, A. L., Nunes, C. M., Fausto, R., & Reva, I. (2018): Conformational control over an aldehyde fragment by selective vibrational excitation of interchangeable remote antennas. Chemical Communications, 54(38), 4778-4781. <https://doi.org/10.1039/C8CC01052H>
- [9] Linden, M. M., Wagner, J. P., Bernhardt, B., Bartlett, M. A., Allen, W. D., & Schreiner, P. R. (2018): Intricate Conformational Tunneling in Carbonic Acid Monomethyl Ester. The journal of physical chemistry letters, 9(7), 1663-1667. <https://doi.org/10.1021/acs.jpcllett.8b00295>
- [10] Marduykov, A., Quanz, H., & Schreiner, P. R. (2017): Conformer-specific hydrogen atom tunnelling in trifluoromethylhydroxycarbene. Nature chemistry, 9(1), 71. <https://doi.org/10.1038/nchem.2609>
- [11] Eckhardt, A. K., Erb, F. R., & Schreiner, P. R. (2019): Conformer-specific [1, 2] H-tunnelling in captodatively-stabilized cyanohydroxycarbene (NC-C [combining umlaut]-OH). Chemical science, 10(3), 802-808. <https://doi.org/10.1039/C8SC03720E>
- [12] Schreiner, P. R., & Reisenauer, H. P. (2008): Spectroscopic identification of dihydroxycarbene. Angewandte Chemie International Edition, 47(37), 7071-7074. <https://doi.org/10.1002/anie.200802105>
- [13] Pettersson, M., Maçôas, E. M., Khriachtchev, L., Fausto, R., & Räsänen, M. (2003): Conformational isomerization of formic acid by vibrational excitation at energies below the torsional barrier. Journal of the American Chemical Society, 125(14), 4058-4059. <https://doi.org/10.1021/ja0295016>

*Last Update:* 2020-01-09 15:05