

# Activation of Small Molecules With Metal-Free Systems

Researchers  
Max Hasenbeck

Principal Investigator  
Dr. Urs Gellrich

Project Term  
2017 - 2020

Clusters  
Goethe-HLR Frankfurt, FUCHS-CSC  
Frankfurt


Software  
GAUSSIAN, ORCA

Institute  
Institute of Organic Chemistry

University  
Justus Liebig University Giessen

Funded by

Fonds der Chemischen Industrie



## Introduction

Ammonia borane (NH<sub>3</sub>BH<sub>3</sub>) is an easy to handle solid capable of releasing up to 3 equivalents of dihydrogen and therefore a potential material for energy storage applications. For the release of dihydrogen at mild conditions accessible and cheap catalysts are highly desirable.

## Methods

Catalyst design and corresponding mechanistic investigations were supported using highlevel quantum mechanical computations.

## Results

We reported the first example of an organocatalyst capable of releasing dihydrogen from ammonia borane at mild conditions which shows comparable activity to noble metal based catalysts (scheme 1).[1] The high level computations helped extensively in elucidating the reaction mechanism of the dehydrogenation of ammonia borane using 6-*tert*-butyl-2-thiopyridone as catalyst.


## Discussion

Computations revealed that the comparable high catalytic activity can be partly attributed to the *tert*-butyl moiety of the catalyst. The catalyst exists in solution in dimeric form. To activate ammonia borane it has to monomerize which is significantly easier when the *tert*-butyl group is present destabilizing the dimeric form because of the steric bulkiness (scheme 2).


## Outlook

Using high level quantum mechanical computations we aim to develop new efficient catalyst for the activation of small molecules either for the release of dihydrogen or the activation of dihydrogen for further chemical transformations.

## Figures



Scheme 1: Dehydrogenation of ammonia borane using 6-*tert*-butyl-2-thiopyridone as catalyst.



Scheme 2: Monomerization of 2-thiopyridone with and without *tert*-butyl group.

## Reference

[1] Hasenbeck, M.; Becker, J.; Gellrich, U.: Efficient Organocatalytic Dehydrogenation of Ammonia Borane, *Angewandte Chemie International Edition* 2020, 59, P.1590-1594. <https://doi.org/10.1002/anie.201910636>

**Last Update:** 2022-04-25 17:46