
Hessisches Kompetenzzentrum
für Hochleistungsrechnen

Intermolecular London Dispersion Interactions of Azobenzene Switches



1 a) X = H,H
b) X = $(CH_2)_2$
c) X = $(CH_2)_3$
d) X = $(CH_2)_4$

Fig. 1: Bis-azobenzenes 1 with different connectors.

Introduction

Azobenzenes (AB) are promising candidates for molecular solar thermal energy storage (MOST) systems. After synthesizing different bisazobenzenes **1 a-d** with different alkyl-substitution on their bridgehead, differential scanning calorimetry (DSC) measurements were performed. It revealed, that intermolecular attractive London dispersion interactions stabilize the *(E)*-isomer in bisazobenzene, due to the different alkyl bridges leading to a higher storage enthalpy. This computational study was performed to support the obtained experimental data. The focus was laid on the storage enthalpies between the all-*(Z)*- and *(E)*-isomers.

Methods

For this project Gaussian16 was used to conduct our computations. All structures were optimized at the PBE0 density functional theory level with D3(BJ) dispersion correction using the def2-TZVP basis set. Molecular structures were visualized using the CYLview software.

Results

By combining the experimental and computational data it was suggested, that in the all-*(E)*-isomer of compounds **1 a-d** there is an intermolecular attractive interaction of the azobenzene with the cycloalkane bridge due to London dispersion. Even though the AB moieties in all molecules investigated are electronically equal increasing London dispersionstrength by different bridging units increased the energy difference between *(E)*- and *(Z)*-state.

Project Manager
Prof. Dr. Hermann A. Wegner

Researchers
Andreas H. Heindl and Anne Kunz

Principal Investigator
Prof. Dr. Hermann A. Wegner

Project Term
2019 - 2020

Clusters
Skylla Cluster Gießen

Software
GAUSSIAN

Institute
Institute of Organic Chemistry

University
Justus Liebig University Giessen

Partners
Department of Chemistry and Chemical Engineering, Chalmers University of Technology (Gothenburg, Sweden)

Funded by

Deutsche Forschungsgemeinschaft

JUSTUS-LIEBIG-UNIVERSITÄT GIESSEN · TECHNISCHE UNIVERSITÄT DARMSTADT · UNIKASSEL VERSITÄT · Philipps Universität Marburg

Those interactions will serve as a basis to design future MOST systems.

Outlook

Future investigations will also be conducted with support from accompanying computational studies on the Skylla Cluster Gießen.

Publications

A. Kunz, A. H. Heindl, A. Dreos, Z. Wang, K. Moth-Poulsen, J. Becker, H. A. Wegner, ChemPlusChem 2019, 84, 1145. https://doi.org/10.1002/cplu.201900330

A. Heindl, L. Schweighauser, C. Logemann, H. Wegner, Synthesis 2017, 49, 2632. https://doi.org/10.1055/s-0036-1589006

Last Update: 2022-06-14 16:32