

# Design and Synthesis of Phosphinidene Chalcogenides

Researchers  
Dr. Artur Mardyukov

Principal Investigator  
Prof. Dr. Peter R. Schreiner

Project Term  
2020 - 2021

Clusters  
Skylia Cluster Gießen, Goethe-HLR  
Frankfurt


Additional Software  
Linux, Windows

Institute  
Institute of Organic Chemistry

University  
Justus Liebig University Giessen

Funded by

Volkswagenstiftung



## Introduction

Phosphinidene chalcogenides (R-P=E; Ch = O, S, Se) are phosphorus analogues of organic nitroso compounds (R-N=O). In contrast to the robust stability of R-N=O, phosphinidene chalcogenides have been considered extremely reactive species.[1-2] These transient species are very reactive toward organic molecules and have been thus used as efficient synthons for new organophosphorus derivatives.[3-5]

Stabilization of R-P=E species can be achieved through coordination to metal centers (thermodynamic stability) or using bulky substituents (kinetic stability). Phenylphosphinidene oxide Ph-P=O (**1**) (the phosphorus analogue of nitrosobenzene) and phenylthiooxophosphine Ph-P=S (**2**), are hitherto unknown experimentally, although the structures and energies of Ph-P=E (E = O, S) and their metal complexes have been computationally and experimentally studied.[6]

## Discussion



Herein, we aim at synthesizing and isolating yet unreported Ph-P=O and Ph-P=S through either flash vacuum pyrolysis or photolysis of their respective azide precursors PhP(E)(N<sub>3</sub>)<sub>2</sub> (E = O, S), thus enabling further studies on their structures and reactivities as ligand-free species. We plan to extend our method to exploit mono-substituted azide precursors. Similar azide architectures have allowed a facile generation of transient nitrenes upon irradiation or thermolysis.[7]

## Outlook

We will study their chemical and physical properties using both experimental and computational methods. Density functional

theory (DFT) and *ab initio* computations will be used to determine the substituent effects on the vibrational as well as optical spectra, photochemistry, geometries, and transition states for subsequent reactions.

## Figures



## Reference

- [1] C. M. E. Graham, J. Valjus, T. E. Pritchard, P. D. Boyle, H. M. Tuononen, P. J. Ragogna, *Inorg. Chem.* 2017, 56, 13500-13509. <https://doi.org/10.1021/acs.inorgchem.7b02217>
- [2] L. Wang, R. Ganguly, F. Mathey, *Organometallics* 2014, 33, 5614-5617. <https://doi.org/10.1021/om500828n>
- [3] C. M. E. Graham, C. L. B. Macdonald, P. D. Boyle, J. A. Wisner, P. J. Ragogna, *Chem. - Eur. J.* 2018, 24, 743-749. <https://doi.org/10.1002/chem.201705198>
- [4] C. M. E. Graham, T. E. Pritchard, P. D. Boyle, J. Valjus, H. M. Tuononen, P. J. Ragogna, *Angew. Chem., Int. Ed.* 2017, 56, 6236-6240. <https://doi.org/10.1002/anie.201611196>
- [5] C. Compain, B. Donnadieu, F. Mathey, *Organometallics* 2005, 24, 1762-1765. <https://doi.org/10.1021/om049089y>
- [6] B. T. Sterenberg, L. Scoles, A. J. Carty, *Coord. Chem. Rev.* 2002, 231, 183-197. [https://doi.org/10.1016/S0010-8545\(02\)00119-4](https://doi.org/10.1016/S0010-8545(02)00119-4)
- [7] X. Zeng, M. Gerken, H. Beckers, H. Willner, *Inorg. Chem.* 2010, 49, 3002-3010. <https://doi.org/10.1021/jc902524u>

*Last Update:* 2022-08-04 16:27