Hessisches Kompetenzzentrum
für Hochleistungsrechnen

Design and Synthesis of Phosphinidene Chalcogenides

Researchers
Dr. Artur Mardyukov

Principal Investigator
Prof. Dr. Peter R. Schreiner

Project Term
2020 - 2021

Clusters
Skylla Cluster Gießen, Goethe-HLR Frankfurt


Additional Software
Linux, Windows

Institute
Institute of Organic Chemistry

University
Justus Liebig University Giessen

Funded by

Volkswagenstiftung



Introduction

Phosphinidene chalcogenides (R–P=E; Ch = O, S, Se) are phosphorus analogues of organic nitroso compounds (R–N=O). In contrast to the robust stability of R–N=O, phosphinidene chalcogenides have been considered extremely reactive species.[1-2] These transient species are very reactive toward organic molecules and have been thus used as efficient synthons for new organophosphorus derivatives.[3-5]

Stabilization of R–P=E species can be achieved through coordination to metal centers (thermodynamic stability) or using bulky substituents (kinetic stability). Phenylphosphinidene oxide Ph–P=O (**1**) (the phosphorus analogue of nitrosobenzene) and phenylthiooxophosphine Ph–P=S (**2**), are hitherto unknown experimentally, although the structures and energies of Ph–P=E (E = O, S) and their metal complexes have been computationally and experimentally studied.[6]

Discussion

Herein, we aim at synthesizing and isolating yet unreported Ph–P=O and Ph–P=S through either flash vacuum pyrolysis or photolysis of their respective azide precursors PhP(E)(N3)2 (E = O, S), thus enabling further studies on their structures and reactivities as ligand-free species. We plan to extend our method to exploit mono-substituted azide precursors. Similar azide architectures have allowed a facilegeneration of transient nitrenes upon irradiation or themolysis.[7]


Outlook

We will study their chemical and physicalproperties using both experimental and computational methods. Density functional


JUSTUS-LIEBIG-UNIVERSITÄT GIESSEN
TECHNISCHE UNIVERSITÄT DARMSTADT
UNIKASSEL VERSITÄT
Philipps Universität Marburg

theory (DFT) and *ab initio* computations will be used to determine the substituent effects on the vibrational as well as optical spectra, photochemistry, geometries, and transitions states for subsequent reactions.

Figures



3, 4 **1, 2**



JUSTUS-LIEBIG-UNIVERSITÄT GIESSEN
TECHNISCHE UNIVERSITÄT DARMSTADT
UNIKASSEL VERSITÄT
Philipps Universität Marburg

Reference

[1] C. M. E. Graham, J. Valjus, T. E. Pritchard, P. D. Boyle, H. M. Tuononen, P. J. Ragogna, Inorg. Chem. 2017, 56, 13500-13509. https://doi.org/10.1021/acs.inorgchem.7b02217

[2] L. Wang, R. Ganguly, F. Mathey, Organometallics 2014, 33, 5614-5617. https://doi.org/10.1021/om500828n

[3] C. M. E. Graham, C. L. B. Macdonald, P. D. Boyle, J. A. Wisner, P. J. Ragogna, Chem. - Eur. J. 2018, 24, 743-749. https://doi.org/10.1002/chem.201705198

[4] C. M. E. Graham, T. E. Pritchard, P. D. Boyle, J. Valjus, H. M. Tuononen, P. J. Ragogna, Angew. Chem., Int. Ed. 2017, 56, 6236-6240. https://doi.org/10.1002/anie.201611196

[5] C. Compain, B. Donnadieu, F. Mathey, Organometallics 2005, 24, 1762-1765. https://doi.org/10.1021/om049089y

[6] B. T. Sterenberg, L. Scoles, A. J. Carty, Coord. Chem. Rev. 2002, 231, 183-197. https://doi.org/10.1016/S0010-8545(02)00119-4

[7] X. Zeng, M. Gerken, H. Beckers, H. Willner, Inorg. Chem. 2010, 49, 3002-3010. https://doi.org/10.1021/ic902524u

Last Update: 2022-08-04 16:27