

Matrix Elements of New Families of Chiral Interactions

Project Manager
Tobias Wolfgruber


Principal Investigator
Prof. Dr. Robert Roth

Project Term
2021 - 2022

Clusters
Lichtenberg Cluster Darmstadt

Institute
Institut für Kernphysik

University
Technische Universität Darmstadt



Introduction

The goal of *ab initio* nuclear structure theory is the description of nuclei from first principles without uncontrolled approximations, allowing calculations with excellent predictive power. All of these calculations require the construction of nuclear interactions, which is done using chiral effective field theory (chiral EFT), a framework that has been extremely successful in recent years. Using these interactions directly in a many-body calculation is computationally expensive, hence, we use the Similarity Renormalization Group (SRG) to ‘soften’ these interactions. The SRG improves the convergence behavior of many-body calculations, however, the SRG itself involves large-scale numerical calculations. This project provides the next generation of chiral interactions suitable for *ab initio* many-body calculations.

Methods

Our main method is the SRG, which is a continuous unitary transformation of the Hamiltonian. This transformation can be written in a flow equation, that translates into a set of coupled nonlinear first-order differential equations. These are solved numerically with highly specialised codes developed within our group. The SRG decouples high- and low-momentum states and thereby prediagonalizes the Hamiltonian, which reduces the model space necessary to converge subsequent many-body calculations.

Results

In this project we have calculated matrix elements of the next generation of chiral interactions, which we developed within the

Low Energy Nuclear Physics International Collaboration (LENPIC) as well as in collaboration with Machleidt. These matrix elements have been used in many large-scale calculations within our group or by our collaborators. Furthermore, we optimized the phenomenological Gaussian four-body interaction and adapted it for tetra-neutron calculations.

Discussion

The systematic order-by-order analysis of the convergence of the chiral EFT expansion and the comparison of different regularization schemes are essential for a quantification of theory uncertainties for nuclear structure observables. With the newest generation of the Entem-Machleidt- Nosyk nucleon-nucleon interaction family in combination with three-nucleon interactions developed in this project, we have an interaction that can provide reliable nuclear observables up into the medium-mass regime.

First discrepancies between theory and experiment have been found in selected open-shell nuclei in the medium mass regime. In the next project period we want to tackle these discrepancies by incorporating the SRG into the fitting procedure, which should enable the construction of even softer interactions.

Furthermore, the consistent SRG evolution of the hyperon-nucleon-nucleon interaction will be performed. Moreover, the development of the interaction family within LENPIC indicates that we are missing crucial contributions either in the interaction or in the radius operator itself. In ongoing work, we will address both higher-order terms of the interaction and the radius operator.

Figures




Figure 1: Ground-state energies and radii of selected medium mass nuclei calculated with the SMS NN interaction supplemented with 3N interactions with $\Lambda=500$ MeV cutoff differing by regularization scheme. The different symbols indicate the c_D -values of the 3N interaction. The results without initial 3N interaction are indicated with open black circles and the experimental values are indicated with black bars.




Figure 2: The low-energy constants c_D and c_E of the nonlocal 3N interaction at N3LO with cutoff 500 MeV optimized for the ground-state energy of ^3H (blue), the ground-state energy of ^4He (red), and the rms relative deviation of the ground-state energies from ^3H and ^4He experiments equal to 0.010 and 0.015 (green areas) and the optimum (green dot). The blue cross and the red square indicate the c_D - c_E pairs that also reproduce the ^{16}O ground-state energy.

Publications

Maris, P.; Epelbaum, E.; Furnstahl, R. J.; Golak, J.; Hebeler, K.; H  ther, T.; Kamada, H.; Krebs, H.; Meißner, Ulf-G.; Melendez, J. A.; Nogga, A.; Reinert, P.; Roth, R.; Skibiński, R.; Soloviiov, V.; Topolnicki, K.; Vary, J. P.; Volkotrub, Yu.; Witała, H.; Wolfgruber, T. (LENPIC Collaboration): Light nuclei with semilocal momentum-space regularized chiral interactions up to third order. Phys. Rev. C 103, 054001 (2021).
<https://doi.org/10.1103/PhysRevC.103.054001>

Reference

Friman-Gayer, U.; Romig, C.; H  ther, T.; Albe, K.; Bacca, S.; Beck, T.; Berger, M.; Birkhan, J.; Hebeler, K.; Hernandez, O. J.; K  nig, S.; Pietralla, N.; Ries, P. C.; Rohrer, J.; Roth, R.; Savran, D.; Scheck, M.; Schwenk, A.; Seutin, R.; Werner, V.: Role of chiral two-body currents in 6-Li magnetic properties in light of a new precision measurement with the relative self-absorption technique. Phys. Rev. Lett. 126, 102501 (2021).
<https://doi.org/10.1103/PhysRevLett.126.102501>

Heil, S.; Petri, M.; Vobig, K.; Bazin, D.; Belarge, J.; Bender, P.; Brown, B. A.; Elder, R.; Elman, B.; Gade, A.; Haylett, T.; Holt, J. D.; H  ther, T.; Hufnagel, A.; Iwasaki, H.; Kobayashi, N.; Loelius, C.; Longfellow, B.; Lunderberg, E.; Mathy, M.; Menendez, J.; Paschalis, S.; Roth, R.; Schwenk, A.; Simonis, J.; Syndikus, I.; Weisshaar, D.; Whitmore, K.: Electromagnetic properties of 21-O for benchmarking nuclear Hamiltonians. Phys. Lett. B 809, 135678 (2020). <https://doi.org/10.1016/j.physletb.2020.135678>

H  ther, T.; Vobig, K.; Hebeler, K.; Machleidt, R.; Roth, R.: Family of Chiral Two- plus Three-Nucleon Interactions for Accurate Nuclear Structure Studies. Phys. Lett. B 808, 135651 (2020).
<https://doi.org/10.1016/j.physletb.2020.135651>

Last Update: 2022-04-23 15:50