
Hessisches Kompetenzzentrum
für Hochleistungsrechnen

# *Theoretical Investigation of Electrocyclic Reactions of Siloles*




## Introduction

Siloles are five-membered heterocyclic dienes, in which a silicon atom replaces one of the carbon atoms of cyclopentadiene. Silole derivatives find application as building units for σ- and π-conjugated compounds for use in organic electroluminescent devices,[1] and their synthesis, reactions, and properties have been subject to intensive studies.[2]

## Results

In an earlier investigation, the 1-chlorosilacyclopentene 1 has been obtained starting from perchlorinated silanes, Li[SitBu3], and 2,3-dimethylbutadiene[3] and siloles can readily be obtained from this compound by deprotonation and LiCl elimination. In a recent combined experimental and theoretical study we have shown that the resulting siloles are very reactive and can isomerise, dimerise or be trapped by reagents like cyclohexene or 2,3-dimethylbutadiene, and the mechanisms underlying these reactions have been investigated in detail by means of double-hybrid density functional theory calculations.[4] Silole 2 can easily isomerise to the more stable silole 3 (Figure 1) via hydrogen shift. Alternatively, it can also dimerise to yield several products: two dimers are formed via Diels-Alder reactions (2+3 and 2+2) and the last one is an head-to-head dimer of 2 and its isomer 3. This latter, more unusual product is accessible via both singlet and triplet intermediates (Figure 1). Further, compound 2 can be trapped prior to isomerisation or dimerisation by addition of excess cyclohexene yielding the racemate of the [4+2] cycloadduct. Under similar conditions, the reaction of 2 with 2,3-dimethylbutadiene yields the [2+4] and [4+2] cycloadducts and hence, 2 acts as silene (dienophile) as well as diene. Our results show that both reaction steps occur without activation barriers (Figure 2).


Researchers
Timo Porsch, Konstantin Falahati and Josef H. Wender

Principal Investigator
Prof. Dr. Max C. Holthausen


Project Term
2015 - 2015

Project Areas
Analytical Chemistry, Method Development (Chemistry)

Clusters
LOEWE CSC Cluster Frankfurt


Institute
Computional Chemistry

University
Goethe Universität Frankfurt am Main





JUSTUS-LIEBIG-UNIVERSITÄT GIESSEN
TECHNISCHE UNIVERSITÄT DARMSTADT
UNIKASSEL VERSITÄT
Philipps Universität Marburg

# Figures





Fig. 1: Formation of silole 2 (and subsequent isomerisation to silole 3) which can dimerise to three different products.





Fig. 2: Barrierless Diels-Alder reaction of the silole (red) with 2,3-dimethylbutadiene can lead to two different products.

# Reference


[1] S. Yamaguchi and K. Tamao (1998): Silole-containing σ- and π-conjugated compounds. J. Chem. Soc. Dalton Trans. 22: 3693-3702. https://doi.org/10.1039/A804491K

[2] J. Dubac, A. Laporterie, and G. Manuel (1990): Group 14 metalloles. 1. Synthesis, organic chemistry, and physicochemical data. Chem. Rev., 1 (90): 215–263. https://doi.org/10.1021/cr00099a008

[3] F. Meyer-Wegner, A. Nadj, M. Bolte, N. Auner, M. Wagner, M.C. Holthausen, and H.-W. Lerner (2011): The Perchlorinated Silanes Si2Cl6 and Si3Cl8 as Sources of SiCl2. Chem. Eur. J., 17 (17): 4715–4719. https://doi.org/10.1002/chem.201003654

[4] F. Meyer-Wegner, J.H. Wender, K. Falahati, T. Porsch, T. Sinke, M. Bolte, M. Wagner, M.C. Holthausen, and H.-W. Lerner (2014): Electrocyclic Reactions of Siloles: A Combined Experimental and Theoretical Study. Chem. Eur. J., 16 (20): 4681–4690. https://doi.org/10.1002/chem.201302544


## *Last Update:* 2022-07-07 15:34