

Density Functional Theory: Study on the Mechanism of the Chloride-Induced Aufbau of Perchlorinated Cyclohexasilanes

Researchers
Lioba Meyer and Dr. Julia I. Schweizer

Principal Investigator
Prof. Dr. Max C. Holthausen

Project Term
2011 - 2014

Project Areas
Analytical Chemistry, Method Development (Chemistry)

Clusters
LOEWE CSC Cluster Frankfurt

Institute
Institut für Anorganische und Analytische Chemie

University
Goethe Universität Frankfurt am Main




Fig. 1: Chloride induced formation of $[\text{Si}_6\text{Cl}_{12}\cdot 2\text{Cl}]^{2-}$.

Introduction

Nano- and micrometer-scaled silicon structures are essential components not only in microelectronic but also in photovoltaic and opto-electronic applications. Oligo- and polymeric (perchloro)silanes are suitable precursors for the formation of silicon wires or thin silicon films.

Methods

The mechanism of the amine-induced disproportionation of perchlorinated silanes affording neo- $\text{Si}_5\text{Cl}_{12}$ was recently established.[1,2] A surprisingly simple preparative procedure, the addition of Si_2Cl_6 to a solution of $[\text{nBu}_4\text{N}]\text{Cl}$ in dichloromethane, leads to the formation of a chloride-complexed cyclic dianion $[\text{Si}_6\text{Cl}_{12}\cdot 2\text{Cl}]^{2-}$ and a variety of its silyl-substituted structural analogs, depending on the reaction conditions. The underlying reaction mechanism has been elucidated by DFT calculations (Fig. 1).

Results

It reveals the chloride ion itself as a Lewis base to trigger a disproportionation of perchlorinated silanes with a subsequent buildup of dianions containing up to eight silicon atoms.[3]

Outlook

The mechanistic insights gained provide the fundament required for the targeted synthesis of oligosilane precursors suitable for microelectronic applications.

Reference

[1] F. Meyer-Wegner, A. Nadj, M. Bolte, N. Auner, M. Wagner, M.C. Holthausen, and H.-W. Lerner (2011), The Perchlorinated Silanes Si_2Cl_6 and Si_3Cl_8 as Sources of SiCl_2 , Chem.-Eur. J. 17, 4715-4719. <https://doi.org/10.1002/chem.201003654>

[2] J. Tillmann, F. Meyer-Wegner, A. Nadj, J. Becker-Baldus, T. Sinke, M. Bolte, M. C. Holthausen, M. Wagner, and H. W. Lerner (2012), Unexpected Disproportionation of Tetramethylethylenediamine-Supported Perchlorodisilane $\text{Cl}_3\text{Si-SiCl}_3$, Inorg. Chem. 51, 8 <https://doi.org/10.1021/ic301283m>

[3] J. Tillmann, L. Meyer, J.I. Schweizer, M. Bolte, H.W. Lerner, M. Wagner, and M.C. Holthausen (2014), Chloride-Induced Aufbau of Perchlorinated Cyclohexasilanes from Si_2Cl_6 : A Mechanistic Scenario, Chem.-Eur. J. 20, 9234-9239. <https://doi.org/10.1002/chem.201402655>

Last Update: 2022-08-12 11:34