

# Interatomic Potentials for Li-Si-O and Si-O-C

Project Manager  
Linus Erhard

Researchers  
Dr. Jochen Rohrer, Marie J. Perera  
(née Charrier), Niklas Leimeroth, Dr.  
Olena Lenchuk and Jara Müller  
Kästner

Principal Investigator  
Prof. Dr. rer. nat. Karsten Albe

Project Term  
2022 - 2023


Clusters  
Lichtenberg II Cluster Darmstadt

Software  
LAMMPS, VASP

Additional Software  
pacemaker

Institute  
Fachgebiet Materialmodellierung

University  
Technische Universität Darmstadt



## Introduction

To understand and study complex materials at the atomic level, it is essential to be able to calculate forces and energies for their atomic configurations. Density functional theory can be used to accurately calculate these properties, but these calculations are computationally expensive and subject to time and length constraints. Machine learning potentials overcome this problem and make it possible to simulate systems with millions of atoms for nanoseconds. In this project, we were interested in comparing several of these machine learning interatomic potentials. In addition, we analysed the highly complex SiO system using an existing machine learning potential and were interested in extending this machine learning potential to other systems. Powerful computing resources are essential for the high-throughput density functional theory calculations for the training data as well as the large-scale molecular dynamics simulation with the machine-learning interatomic potentials.

## Methods

As input data for the machine learning potentials, we use training data labelled with energies and forces from density

functional theory. There are different types of machine learning potentials, depending on the machine learning method used and the descriptor of the atomic environment. In our case, we tested the Gaussian approximation potentials, the moment tensor potential and the atomic cluster expansion.

## Results

### 1. Benchmarking of different machine-learning interatomic potential approaches

For atomistic simulations, it is essential that the interatomic potential used is computationally efficient and accurate. There are now many types of machine-learning interatomic potential available. To find out which of these potentials offers the best performance for our system, we compare atomic cluster expansion, moment tensor potentials and Gaussian approximation potentials. We found that for the SiO system, the non-linear Atomic Cluster Expansion provides the best trade-off between accuracy and speed.

### 2. Generation of large scale silicon monoxide structures

The structure of silicon monoxide has long been controversial. Experiments have shown that it is a mixture of amorphous silicon and silicon dioxide. However, with only nanometre-sized grains. Until now, an atomistic description has not been possible because an exact description of the forces and energies on the required length scales was missing. We are now able to use a machine learning interatomic potentials to create structural models of silicon monoxide that are in excellent agreement with the experiment (see Fig. 1).

### 3. Construction of a Si-O-C database

Silicon oxycarbides are interesting for a wide range of applications. They can be used for structural applications as well as battery material in lithium batteries. We have extended the existing potential for SiO to carbon and significantly increased the database of density functional theory data. With this new potential, we were able to create structures containing phase-separated carbon and silicon dioxide as well as silicon carbide (see Fig. 2).

### 4. Construction of a Li-Si-O database

Unfortunately, due to time constraints, it was not possible to complete this part of the project within the project duration

## Discussion

Through machine learning interatomic potentials trained on a large amount of density functional theory data, we can gain new insights into materials. We have demonstrated this by generating realistic nanostructures of SiO and SiOC. In the future, these machinelearning interatomic potentials could be used to investigate a wide range of other problems in SiO and SiOC to improve our understanding of these systems. Future work is also planned to extend the machine learning interatomic potential models to lithium.

## Figures




Figure 1: Atomistic model of silicon monoxide generated by an ACE potential for SiO.




Figure 2: SiOC structural model created by a melt-quench simulation using the ACE potential for SiOC.

## Publications

Erhard, L. C.; Rohrer, J.; Albe, K.; Deringer, V. L.: "Modelling atomic and nanoscale structure in the silicon-oxygen system through active machine learning", Nat Commun 15, 1927 (2024)  
<https://doi.org/10.1038/s41467-024-45840-9>

Leimeroth, N.; Rohrer, J.; Albe, K.: "Structure-Property relations of silicon oxycarbides studied using a novel machine learning interatomic potential", in preparation

Erhard, L.C.; Leimeroth, N.; Rohrer, J.; Albe, K.: "Benchmarking of various machine-learning interatomic potentials for SiO and AlCuZr", in preparation

Linus Erhard, Benchmarking of different machine-learning interatomic potential approaches for silica (oral), MMM10, Baltimore, US, October 2-7, 2022

Linus Erhard, A machine-learned interatomic potential for silica and mixed silica-silicon systems (oral), DPG Spring Meeting, Dresden, Germany, March 26-31, 2023

Linus Erhard, Modelling atomic and nanoscale structure in the silicon-oxygen system through active machine learning (oral), GAP/(m)ACE Developers and Users Meeting, Coventry, UK, September 12-15, 2023

Linus Erhard, Unraveling the atomic structure of silicon monoxide using high-performance computing and machine learning (oral), NHR Conference, Berlin, Germany, September 18-19, 2023

*Last Update:* 2024-12-20 12:22