



Hessisches Kompetenzzentrum
für Hochleistungsrechnen

Dynamical Simulation of Heavy Ion Reactions at FAIR

Researchers

Jan Steinheimer, D. Oliinychenko
and J. Auvinen

Principal Investigator

Prof. Dr. Hannah Petersen

Project Term

2015 - 2015

Project Areas

Optics, Quantum Optics and Physics
of Atoms, Molecules and Plasmas

Clusters


LOEWE CSC Cluster Frankfurt

Institute

Institut für Theoretische Physik

University

Goethe Universität Frankfurt am
Main



Introduction

In heavy ion collisions at lower beam energies the QCD phase diagram is probed at high net baryon densities. To gain insights about the equation of state of strongly interacting matter and the transport properties of the newly produced state of matter, the quark gluon plasma, sophisticated dynamical models are needed.

Methods

In this work a hybrid approach based on ideal relativistic fluid dynamics for the hot and dense stage and hadronic transport for the initial and final non-equilibrium stage is used to explore flow observables in a wide range of beam energies, that has been recently investigated experimentally in the beam energy scan program at the Relativistic Heavy Ion Collider at Brookhaven National Laboratory.

This kind of dynamical simulations requires high performance computing since event by event calculations are essential to compare to experimental data in a realistic fashion. Anisotropic flow can be quantified by the Fourier coefficients of the azimuthal distribution of the produced particles and is particularly sensitive to the transport properties, such as the

shear viscosity, of the produced system. At high collision energies it has been one of the main signals indicating that the quark gluon plasma behaves as a perfect liquid.

Results

Therefore, it was rather surprising that the amount of elliptic flow is rather insensitive to a change in beam energy. Our calculations (see Fig. 1) show that v_2 produced by hydrodynamics is considerably diminished at lower collision energies, but this decrease is partially compensated by the transport dynamics[1,2]. The pre-hydrodynamics transport phase is of particular importance for understanding the collision energy evolution, while the hadronic rescatterings after the hydrodynamical phase contribute more systematically $\sim 10\%$ to the total flow at all energies. Recent STAR data from the RHIC beam energy scan show that the slope of the net-proton directed flow v_1 near midrapidity changes sign twice within the investigated collision energy range which has been predicted by earlier fluid calculations to be a signal of a first-order phase transition between hadronic matter and quarkgluon plasma. We examine the sensitivity of the directed flow v_1 to the order of the phase transition by comparing hybrid simulations with a first-order phase transition „Bag model“ equation of state (EoS) to calculations with a chiral model EoS, which has a cross-over phase transition. As can be seen in Fig. 2 the two EoS are completely indistinguishable in the hybrid simulations, questioning the usability of v_1 as a signal of the first-order phase transition[3,4].

Outlook

However, the hybrid model results also deviate notably from experimental data, which makes further studies on the topic necessary.

Figures




Fig. 1: Magnitude of v_2 in midcentral collisions at the beginning of hydrodynamical evolution (diamonds), immediately after the end of hydrodynamics phase (squares) and after the full simulation (circles).




Fig. 2 : v_1 slope at midrapidity for protons (solid symbols) and anti-protons (open symbols) for impact parameter range, extracted from the hybrid model calculations with a first order (black) and crossover EoS (red). Results are compared with the UrQMD model calculations (grey) and experimental data from STAR, NA49 and E895 collaborations (green).

Reference

1. J. Auvinen and H. Petersen (2013), Collision Energy Evolution of Elliptic and Triangular Flow in a Hybrid Model, PoS CPOD 2013, 034.
2. J. Auvinen and H. Petersen (2013), Evolution of elliptic and triangular flow as a function of collision energy in a hybrid model, Phys. Rev. C 88, 064908. <https://doi.org/10.1103/PhysRevC.88.064908>
3. J. Steinheimer, J. Auvinen, H. Petersen, M Bleicher, and H. Stöcker (2014), Examination of directed flow as a signal for a phase transition in relativistic nuclear collisions, Phys. Rev. C 89, 054913. <https://doi.org/10.1103/PhysRevC.89.054913>
4. J. Auvinen, J. Steinheimer, and H. Petersen (2014), What the collective flow excitation function can tell about the quarkgluon plasma, Nucl. Phys. A 931, 975. <https://doi.org/10.1016/j.nuclphysa.2014.09.056>

Last Update: 2022-06-30 18:12